

WEST Search History

DATE: Tuesday, December 02, 2003

<u>Set Name</u>	<u>Query</u>	<u>Hit Count</u>	<u>Set Name</u>
side by side			result set
	DB=DWPI; PLUR=YES; OP=ADJ		
L8	(snip or snlp) and l5	0	L8
L7	L5 and proteasome	1	L7
L6	tongwen-wang\$.in. and (smad or smad1)	0	L6
L5	("WANG-T".IN. "WANG-T-T".IN. "WANG-T-S-F".IN. "WANG-T-S".IN. "WANG-T-R".IN. "WANG-T-P".IN. "WANG-T-M".IN. "WANG-T-K".IN. "WANG-T-J".IN. "WANG-T-L".IN. "WANG-T-T-M".IN. "WANG-T-T-T".IN. "WANG-T-W".IN. "WANG-T-W-Y".IN. "WANG-T-X".IN. "WANG-T-Y".IN.)!	1060	L5
L4	tongwen-wang\$.in.	0	L4
L3	wang-tongwen\$.in.	0	L3
L2	pctus0003561	0	L2
	DB=USPT; PLUR=YES; OP=ADJ		
L1	5912224.pn.	1	L1

END OF SEARCH HISTORY

WEST**End of Result Set**

Generate Collection

Print

L1: Entry 1 of 1

File: USPT

Jun 15, 1999

US-PAT-NO: 5912224

DOCUMENT-IDENTIFIER: US 5912224 A

TITLE: Methods and compositions for enhancing cellular response to TGF-.beta. ligands

DATE-ISSUED: June 15, 1999

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Donahoe; Patricia K.	Weston	MA		
Wang; Tongwen	Arlington	MA		

US-CL-CURRENT: 514/2; 424/198.1, 424/85.1, 514/12, 514/291, 530/399

CLAIMS:

We claim:

1. A method for potentiating a cellular response, comprising:

(a) administering to cells which express a receptor of a Transforming Growth Factor-beta (TGF-.beta.) family ligand an effective amount of a TGF-.beta. family ligand to induce the receptor-mediated cellular response; and

(b) administering to said cells an effective amount of a macrolide which is a naturally occurring or synthetic FK506 or rapamycin derivative, lacks or has reduced immunosuppressive activity as compared to FK506 or rapamycin, respectively, and binds FKBP12 to potentiate the cellular response.

2. The method of claim 1, wherein said macrolide is a FK506 antagonist.

3. The method of claim 2, wherein said FK506 antagonist has the following formula: ##STR3## wherein R.sub.1 is selected from the group consisting of H, alkyl, aryl and acyl, R.sub.2 is selected from the group consisting of H and --OR.sub.5, such that R.sub.5 is selected from the group consisting of H, alkyl, aryl and acyl, R.sub.3 is selected from the group consisting of --CH.sub.2 -- and --CH.sub.2 CH.sub.2 --, and R.sub.4 is selected from the group consisting of H, alkyl, acyl and aryl.

4. The method of claim 2, wherein said FK506 antagonist has the following formula: ##STR4## wherein R.sub.1 is selected from the group consisting of H and --CH.sub.3, R.sub.2 is selected from the group consisting of H and --OH, and R.sub.3 is selected from the group consisting of --CH.sub.2 -- and --CH.sub.2 CH.sub.2 --.

5. The method of claim 2, wherein said FK506 antagonist is 15-O-desmethyl-FK520.

6. The method of claim 2, wherein said FK506 antagonist is L-685,818.

7. The method of claim 1, wherein said receptor is selected from the group consisting of R1, R2, R3, R4, ALK3, ALK6, Sax and Tkv.